

Australian Standard™

Coal and coke—Analysis and testing

**Part 10.2: Determination of trace elements—Coal and coke—
Determination of arsenic, antimony and selenium—Hydride generation method**

This Australian Standard was prepared by Committee MN/1, Coal and Coke. It was approved on behalf of the Council of Standards Australia on 27 February 1998 and published on 5 June 1998.

The following interests are represented on Committee MN/1:

ACIRL

Australasian Institute of Mining and Metallurgy

Australian Coal Association

Australian Coal Preparation Society

Australian Institute of Energy

Bureau of Steel Manufacturers of Australia

Coalfield Geology Council of New South Wales

CSIRO, Division of Coal and Energy Technology

Department of Mines and Energy, Queensland

Electricity Supply Association of Australia

Minerals Council of Australia

Institution of Engineers, Australia

Royal Australian Chemical Institute

University of New South Wales

University of Newcastle

University of Queensland

Review of Australian Standards. To keep abreast of progress in industry, Australian Standards are subject to periodic review and are kept up to date by the issue of amendments or new editions as necessary. It is important therefore that Standards users ensure that they are in possession of the latest edition, and any amendments thereto.

Full details of all Australian Standards and related publications will be found in the Standards Australia Catalogue of Publications; this information is supplemented each month by the magazine 'The Australian Standard', which subscribing members receive, and which gives details of new publications, new editions and amendments, and of withdrawn Standards.

Suggestions for improvements to Australian Standards, addressed to the head office of Standards Australia, are welcomed. Notification of any inaccuracy or ambiguity found in an Australian Standard should be made without delay in order that the matter may be investigated and appropriate action taken.

This Standard was issued in draft form for comment as DR 96493.

Australian Standard™

Coal and coke—Analysis and testing

**Part 10.2: Determination of trace elements—Coal and coke—
Determination of arsenic, antimony and selenium—Hydride generation method**

Originated as AS K152.10—1965.
Previous edition AS 1038.10—1980.
Revised and redesignated AS 1038.10.2—1998.

PREFACE

This Standard was prepared by the Standards Australia Committee MN/1, Coal and Coke, as a revision of AS 1038.10, *Coal and Coke—Analysis and testing*, Part 10: *Arsenic in coal and coke*.

The objective of this revision is to include the determination of antimony and selenium and to update the method for determination of arsenic.

The determination of the trace elements in coal and coke is becoming more important due to the considerable emphasis being placed on the effect of these elements on the environment. International buyers are becoming increasingly aware of the need for more detailed knowledge of the coals that they are purchasing and may request trace element analysis. In order to have accurate and precise results for this analysis it is imperative that standard methods be available.

© Copyright — STANDARDS AUSTRALIA

Users of Standards are reminded that copyright subsists in all Standards Australia publications and software. Except where the Copyright Act allows and except where provided for below no publications or software produced by Standards Australia may be reproduced, stored in a retrieval system in any form or transmitted by any means without prior permission in writing from Standards Australia. Permission may be conditional on an appropriate royalty payment. Requests for permission and information on commercial software royalties should be directed to the head office of Standards Australia.

Standards Australia will permit up to 10 percent of the technical content pages of a Standard to be copied for use exclusively in-house by purchasers of the Standard without payment of a royalty or advice to Standards Australia.

Standards Australia will also permit the inclusion of its copyright material in computer software programs for no royalty payment provided such programs are used exclusively in-house by the creators of the programs.

Care should be taken to ensure that material used is from the current edition of the Standard and that it is updated whenever the Standard is amended or revised. The number and date of the Standard should therefore be clearly identified.

The use of material in print form or in computer software programs to be used commercially, with or without payment, or in commercial contracts is subject to the payment of a royalty. This policy may be varied by Standards Australia at any time.

CONTENTS

	<i>Page</i>
1 SCOPE	4
2 REFERENCED DOCUMENTS	4
3 DEFINITIONS	4
4 PRINCIPLE	4
5 SAFETY	4
6 REAGENTS	4
7 APPARATUS	6
8 SAMPLE	6
9 PROCEDURE	6
10 MEASUREMENT BY ATOMIC ABSORPTION OR ATOMIC FLUORESCENCE SPECTROMETRY AFTER HYDRIDE GENERATION	8
11 CALCULATION	8
12 REPORTING OF RESULTS	8
13 PRECISION	8
14 TEST REPORT	9

STANDARDS AUSTRALIA

Australian Standard

Coal and coke—Analysis and testing

Part 10.2: Determination of trace elements—
Coal and coke—Determination of arsenic, antimony and
selenium—Hydride generation method

1 SCOPE This Standard sets out a method for the determination of arsenic, antimony and selenium in higher rank coal and coke by hydride generation.

2 REFERENCED DOCUMENTS The following documents are referred to in this Standard:

AS

1038	Coal and coke—Analysis and testing
1038.3	Part 3: Proximate analysis of higher rank coal
1038.4	Part 4: Coke—Proximate analysis
1038.16	Part 16: Assessment and reporting of results
2134	Recommended practice for chemical analysis by atomic absorption spectrometry
2134.3	Part 3: Vapour generation atomic absorption spectrometry
2243	Safety in laboratories (series)
2418	Coal and coke—Glossary of terms
2508	Safe storage and handling information cards (series)
2706	Numerical values—Rounding and interpretation of limiting values
4264	Coal and coke—Sampling
4264.1	Part 1: Higher rank coal—Sampling procedures
4264.2	Part 2: Coke—Sampling procedures

3 DEFINITIONS For the purpose of this Standard, the definitions given in AS 2418 apply.

4 PRINCIPLE A known mass of sample is ignited in intimate contact with Eschka mixture in an oxidizing atmosphere at 800°C to decompose the organic matter. The residue is then extracted with hydrochloric acid and the analytes are determined by atomic absorption or atomic fluorescence spectrometry after conversion to their volatile hydrides.

5 SAFETY For information on laboratory safety, reference should be made to the relevant parts of AS 2243 and AS 2508.

Reagents used may be hazardous or toxic and reference should be made to the appropriate Material Safety Data Sheets.

6 REAGENTS

6.1 General Unless otherwise specified, all reagents shall be of analytical reagent grade, and only distilled water or water of equivalent purity shall be used.