

1994 ED.

AS 2636—1988



**Standards
Association of
Australia**



Australian Standard[®] 2636—1988

SAMPLING OF VENOUS AND CAPILLARY BLOOD FOR THE DETERMINATION OF LEAD OR CADMIUM CONCENTRATION

AS 2636—1994
Sampling of venous and
capillary blood for the
determination of lead or
cadmium concentration 8pp D
Specifies the procedures for the
sampling of blood, using needle
and syringes or vacuum systems,
for the subsequent determination
of cadmium or lead. The
Standard also includes appendices
which describe the method for
determining the contaminant
analyte level in collection
equipment, containers and swabs.
Supersedes AS 2636—1988. Publication
date 1994-04-11.



This Australian Standard was prepared by Committee CH/6, Analysis of Body Fluids for Metals Content. It was approved on behalf of the Council of the Standards Association of Australia on 21 January 1988 and published on 5 April 1988.

The following interests are represented on Committee CH/6:

Adelaide Children's Hospital Inc.
Australasian Institute of Mining and Metallurgy
Australian Association of Clinical Biochemists
Australian Institute of Medical Laboratory Scientists
Australian Mineral Development Laboratories
Department of Health, N.S.W.
Department of Labour, Vic.
National Health and Medical Research Council
Nutrition Society of Australia
Prince of Wales Hospital, N.S.W.
Royal Australian Chemical Institute
Royal Prince Alfred Hospital, N.S.W.
Royal College of Pathologists of Australia

Review of Australian Standards. To keep abreast of progress in industry, Australian Standards are subject to periodic review and are kept up-to-date by the issue of amendments or new editions as necessary. It is important therefore that Standards users ensure that they are in possession of the latest edition, and any amendments thereto.

Full details of all SAA publications will be found in the Catalogue of SAA Publications; this information is supplemented each month by SAA's journal 'The Australian Standard', which subscribing members receive, and which gives details of new publications, new editions and amendments, and of withdrawn Standards.

Suggestions for improvements to Australian Standards, addressed to the head office of the Association, are welcomed. Notification of any inaccuracy or ambiguity found in an Australian Standard should be made without delay in order that the matter may be investigated and appropriate action taken.

This Standard was issued in draft form for comment as DR 87073.

AUSTRALIAN STANDARD

**SAMPLING OF VENOUS AND
CAPILLARY BLOOD FOR THE
DETERMINATION OF LEAD OR
CADMIUM CONCENTRATION**

AS 2636—1988

First published as AS 2636—1983.
Second edition 1988.

**PUBLISHED BY THE STANDARDS ASSOCIATION OF AUSTRALIA
STANDARDS HOUSE, 80 ARTHUR ST, NORTH SYDNEY, N.S.W.**

ISBN 0 7262 4919 X

PREFACE

This Standard was prepared by the Association's Committee for the Analysis of Body Fluids for Metals Content under the direction of the Chemical Standards Board. This Standard supersedes AS 2636—1983, *Sampling of venous and capillary blood for the determination of lead content*.

The sampling of blood for the determination of lead and cadmium presents a number of problems, the major one being the possibility of contamination of the sample. This Standard has been prepared to provide advice about the best procedure for obtaining samples of both venous and capillary blood, free from contamination and suitable for use in the determination of lead and/or cadmium concentration.

Venous blood samples should be used in preference to capillary samples to minimize contamination with lead or cadmium from the skin or from other sources. Capillary samples are suitable for screening large populations, provided that elevated concentrations of lead or cadmium are confirmed immediately by the analysis of venous samples.

This Standard should be read in conjunction with the analytical methods for lead (AS 2787 and AS 2411) and cadmium (AS 3503).

CONTENTS

	<i>Page</i>
1 SCOPE	3
2 REFERENCED DOCUMENTS	3
3 SAMPLING OF VENOUS BLOOD	3
4 SAMPLING OF CAPILLARY BLOOD.....	3
APPENDIX A. METHOD FOR DETERMINATION OF LEAD OR CADMIUM CONTAMINATION FROM SYRINGES, TUBES AND SWABS	5

© Copyright — STANDARDS ASSOCIATION OF AUSTRALIA 1988

Users of Standards are reminded that copyright subsists in all SAA publications. Except where the Copyright Act otherwise allows, no part of this publication may be reproduced, stored in a retrieval system in any form or transmitted by any means without prior permission in writing of the Standards Association of Australia. Requests for permission should be directed to the Head Office of the Association. Where such requests relate to the reproduction of the whole or a substantial part of any Standard, permission may be conditional on an appropriate royalty payment.

STANDARDS ASSOCIATION OF AUSTRALIA

Australian Standard

SAMPLING OF VENOUS AND CAPILLARY BLOOD FOR THE DETERMINATION OF LEAD OR CADMIUM CONCENTRATION

1 SCOPE. This Standard sets out procedures for the sampling of venous and capillary blood for the subsequent determination of their lead and cadmium concentration. It does not extend to the collection of serum and plasma. A method for the determination of lead or cadmium contamination from syringes, tubes and swabs is set out in Appendix A.

NOTE: In this document analyte refers to lead and/or cadmium.

2 REFERENCED DOCUMENTS. The documents below are referred to in this Standard:

AS

2134 Recommended practice for chemical analysis by atomic absorption spectrometry
Part 1: Flame atomic absorption spectrometry
(AS 2134.1)

2787 Whole blood—Determination of lead—Electro-thermal atomization atomic absorption spectrometric method

3503 Whole blood—Determination of cadmium—Furnace atomic absorption spectrometric method

3 SAMPLING OF VENOUS BLOOD.

3.1 General. Blood is drawn from the vein in the ante-cubital fossa, or any other suitable vein, normally by means of a disposable needle and syringe.

3.2 Equipment.

3.2.1 General. All equipment and reagents used for the handling and treatment of blood samples shall be checked for either lead or cadmium contamination as appropriate. Suitable procedures for performing these checks and maximum contamination levels acceptable are contained in Appendix A.

3.2.2 Plasticsware. Both coloured and non-coloured plasticsware may be used. In some cases however, coloured plasticsware may give rise to contamination from cadmium colouring compounds and should be duly tested by the procedure described in Appendix A.

3.2.3 Syringes and needles. Each batch of syringes and needles used shall be examined by systematic sampling to ascertain whether or not these items contain a significant amount of the analyte.

3.2.4 Blood containers. Blood containers shall be shown to be free of significant amounts of the analyte in accordance with the procedure described in Appendix A. A wide variety of containers is available and the containers are usually supplied with anticoagulants. Suitable anticoagulants are lithium heparin, ammonium heparin, sodium heparin, and di-potassium and di-sodium salts of ethylenediaminetetra-acetic acid (EDTA).

Recommended levels of anticoagulant are—

(a) heparin—10 IU/mL to 15 IU/mL of whole blood;
or

(b) EDTA (di-potassium salt)—1 to 1.5 mg/mL of whole blood.

These materials, whether in the tube at purchase or subsequently added to the container, shall be shown to be free of significant amounts of the analyte.

NOTE: Heparin or any of its preparations is not recommended unless the sample is to be analysed within 48 h. These anticoagulants may lead to the formation of clots upon storage of the sample.

3.2.5 Swabs. Swabs or preparations used in conjunction with swabs, shall be shown to be free of significant amounts of the analyte in accordance with the procedure described in Appendix A.

3.3 Preparation of the sampling site. The procedure shall be as follows:

(a) If the arm is soiled, wash the area around the vein thoroughly with soap and water.

(b) Wipe the site vigorously with an alcohol impregnated swab (4.2.4(a)).

(c) Dry the site using absorbent sterile cotton wool or gauze swab (4.2.4(b)).

3.4 Collection of venous blood. The blood specimen is ideally taken after the patient has been seated for 10 min. The concentration of lead in the blood sample is related to the packed red cell volume, hence the use of a tourniquet should be avoided if possible. Should a tourniquet be required to enable the vein to be located, it is preferable for the tourniquet to be released prior to the withdrawal of the sample.

3.5 Transfer of the sample to the container and storage of the sample. The blood shall be transferred from the syringe as soon as possible to avoid coagulation of the sample in the syringe. Both the container size and the amount of anticoagulant should be matched to the volume of the blood sample collected.

The needle shall be removed from the syringe before the blood is transferred to the sample container. Squirting the blood through the needle can cause the blood to haemolyse, rendering the sample unsuitable for haematological examination.

The blood sample shall be mixed thoroughly by inversion.

If the sample is to be split, it is advisable to introduce air into the syringe after it has been withdrawn from the patient so that several inversions of the syringe will ensure mixing of the blood to avoid any possibility of stratification.

4 SAMPLING OF CAPILLARY BLOOD.

4.1 General. Capillary blood samples are taken by puncturing the skin to obtain a free flow of blood.

Up to 1.0 mL of blood may be collected by this procedure.